

요추부 수술 후 경막외 무통방법과 지속적 정맥 주입에 의한 통증 조절의 효과 비교

오규성 · 신문수 · 길현주* · 허 룡 · 최훈규 · 안정용 · 권성오**

= Abstract =

Effects of Continuous Intravenous Analgesia Versus Epidural Analgesia after Lumbar Spinal Surgery : A Prospective Study

Kyu Sung Oh, M.D., Moon Soo Shin, M.D., Hyun Joo Gil, M.D.,*

Ryoong Hur, M.D., Hun Kyu Choi, M.D.,

Jung Yong Ahn, M.D., Seong Oh Kwon, M.D.**

*Department of Neurosurgery and Anesthesiology, * Pundang CHA Hospital, Pochon CHA University College of Medicine, Sungnam, Korea*

*Department of Neurosurgery, ** Gumi CHA Hospital, Pochon CHA University College of Medicine, Gumi, Korea*

Objective : The purpose of this non - randomized prospective study was to evaluate the safety and efficacy of continuous intravenous nalbuphine - ketorolac - droperidol(CIA) versus continuous infusion of epidural morphine - bupivacaine(CEA) for pain control after lumbar spinal surgery.

Methods : Twenty - one patients who underwent spine surgery including laminectomy, fusion with fixation were assigned to receive an intravenous bolus of nalbuphine 5mg and ketorolac 15mg, followed by a continuous infusion of nalbuphine 25mg, ketorolac 105mg, and droperidol 5mg mixed with normal saline 98cc(2cc/hr). Twenty patients received a bolus infusion of morphine 2mg and 0.125% bupivacaine 8cc followed by a continuous intravenous infusion of 100cc 0.125% bupivacaine and morphine sulfate 8.0mg(2cc/hr). Pain score was measured on a visual analogue scale(VAS). It's safety and efficacies were compared with the results of continuous infusion of epidural morphine - bupivacaine, which was reported previously by same authors. A continuous infuser was used to give epidural morphine - bupivacaine and intravenous nalbuphine - ketorolac - droperidol.

Results : In general, mild pain, pain less than 3 VAS scores, was observed postoperatively from 30minutes to 72hours in CEA group, and from 6 hours to 72 hours in CIA group. The early postoperative pain was controlled easily in 6 hours in CEA group, compared to CIA group($p < 0.05$). However, there was no statistical significance in 72 hours on pain scores between CEA and CIA groups after 6 - 12hours of pain managements. Pruritus, nausea and vomiting, and urinary retention were more frequent in CEA group.

Conclusion : CIA and CEA are considered effective methods in postoperative pain managements. However, adequate doses in early intravenous infusion and continuous intravenous analgesia with nalbuphine - ketorolac - droperidol will be needed for better control in early postoperative pain with less side effects.

KEY WORDS : Epidural anesthesia · Intravenous anesthesia · Nalbuphine · Ketorolac · Droperidol · Lumbar spinal surgery.

서 론

가

가




Fig. 1. Pain scores measured by Visual Analogue Scale in lami-nectomy group. * : $p < 0.05$ compared between CEA group and CIA group.




Fig. 2. Pain scores measured by Visual Analogue Scale in fu-sion group. * : $p < 0.05$ compared between CEA group and CIA group.




Fig 3. Pain scores measured by Visual Analogue Scale in Pa-tient data. * : $p < 0.05$ compared between CEA group and CIA group.

($p > 0.05$)(Fig. 3).

CEA
30 , CIA
VAS 3
CIA
48
VAS 가 3 , CIA
6

Table 1. Side effects due to continuous epidural analgesic

Side effect	Number of patient(%)
Pruritus	1/20(5%)
Nausea/vomiting	2/20(10%)
Urinary retension	2/20(10%)
Respiratory depression	0/20(0%)
Total	4/20(25%)

Table 2. Side effects due to continuous intravenous analgesic

Side effect	Number of patient(%)
Nausea/vomiting	1/22(4.5%)
Respiratory depression	0/22(0%)
Cardio-vascular problem	0/22(0%)
Total	1/22(4.5%)

6~12

가 (p=0.062).
CEA / 가 2
1 , 가 1 (Table 1),
CIA / 가 1
(Table 2).

고 찰

nalbuphine - μ -
K - , 가
morphine 60~80% 2)13)15), morphine
가 . Morphine
1998 CEA
morphine
Caudal block morphine
7.9% 가
19). Nalbuphine
opioid
12)13)
Ketorolac NSAIDs cyclooxygenase
prostaglandin
, ketorolac

morphine morphine , morphine nalbuphine
 ,
 1)6)18) . Opioid CEA CIA
 , ,
 , 48
 (Chemoreceptor trigger zone) 가 (psychotomimetic reaction),
 3)5) . Ketorolac 30 2).
 60mg 15 30mg 6 CIA 가 CEA
 , 10 8). 12
 CEA
 가 , 48
 CEA 가
 nalbuphine CEA
 nalbuphine 5mg keto- 48
 rolac 15mg , 가
 nalbuphine 25mg, ketorolac
 105mg, droperidol 5mg normal saline 90ml 48 , CIA
 100ml가 2ml가 CEA
 가
 , nalbuphine 50mg 21 8 48
 40 4 , 3 , 4 가
 2). morphine CEA CIA
 가 nalbu- 6
 phine 30mg/70kg , 12 , 48
 10). 48 VAS 3
 nalbuphine 25mg 가
 , 1 (rebound phenomenon)
 CIA CIA
 가 12 6 CIA
 nalbuphine , ketorolac , CEA
 가 가 nalbuphine
 가 , opioid , 가
 droperidol 가 . Drope- 4).
 ridol butyrophenone dopamin
 ,
 CEA 2 , CIA
 1 가
 가
 . CIA CEA

결 론

6
가 , 가
, CEA 가
CEA
6
nalbuphine CIA
가
.
• : 2001 7 10
• : 2001 10 29
• :
463 - 712 351
: 031) 780 - 5260, : 031) 780 - 5269
E - mail : smsnsd2000@yahoo.co.kr

References

- 1) Ahn HJ, Kim MH : *The effects of ketorolac on T cell subsets in patients undergoing a gastrectomy. J Kor Anaesth Soc* 40 : 201-209, 2001
- 2) Bang EC, Kim SY, Lee HS, Kang YI, Kim MH, Cho KS : *The presumptive analgesia with intravenous nalbuphine-ketorolac in gynecologic surgery. J Kor Pain Soc* 13 : 38-43, 2000
- 3) Bollish SJ, Collins CL, Kirling DM, Bartlett RH : *Efficacy of patient controlled versus conventional analgesia for postoperative pain. Clin Pharm* 4 : 48-52, 1985
- 4) Cho HS, Lee JU : *Analgesic effect of iv-PCA with nalbuphine-ketorolac for upper abdominal surgical patients. J Kor Anaesth Soc* 37 : 86-91, 1997
- 5) Dahl JB, Dagaard JJ, Larsen HV : *Patient-controlled analgesia : A controlled trial. Acta Anaesthesiol Scand* 31 : 744-777, 1987
- 6) Gillies GWA, Kenny GNC, Bullingham RES, McArdle CS : *The morphine sparing effects of ketorolac tromethamine : A study of a new parenteral non-steroidal anti-inflammatory agent after abdominal surgery. Anaesthesia* 42 : 727-731, 1987
- 7) Han JS, Park SK, Suh MS, Yoo ES, Lee SY, Min BH, et al : *Effects of Intravenous and epidural patient-controlled analgesia in prospective pain and knee rehabilitation after a unilateral total knee replacement. J Kor Anaesth Soc* 40 : 47-53, 2001
- 8) Kenny GNC : *Ketorolac trometamol : A new non-opioid analgesic. Br J Anaesth* 65 : 445-447, 1990
- 9) Kim ST, Jun JH, Jeon JW, Kim DW, Shim JC, Kim KH, et al : *Comparison of an effective dose of intravenous postoperative patient-controlled analgesia with nalbuphine. J Kor Anaesth Soc* 40 : 195-200, 2001
- 10) Lee JS, Lee YW, Yoon DM, Nam YT, Song KH : *Intravenous patient-controlled analgesia with nalbuphine : could be an alternative to epidural patient-controlled analgesia with morphine-bupivacaine for pain relief after cesarian delivery?. J Kor Anaesth Soc* 10 : 34-39, 1997
- 11) Lehmann KA : *Update of patient-controlled analgesia. Curr Opin Anaesthesiol* 10 : 374-379, 1996
- 12) Miller RD : *Anesthesia. 5th ed. New York. Churchill Livingstone, 2000, pp343-347*
- 13) Miller RR : *Evaluation of nalbuphine hydrochloride. Am Hosp Pharm* 37 : 942-9, 1980
- 14) Niv D, Wolman I, Alon E, Weinbroum A, Varrassi G : *Morphine versus nalbuphine for postoperative pain relief using the patient-controlled analgesia method. Eur J Pain* 16 : 1-2, 1995
- 15) Pugh G, Drummond G : *A dose-response study with nalbuphine hydrochloride for pain in patients after upper abdominal surgery. Br J Anaesth* 59 : 1356-1363, 1987
- 16) Romagnoli A, Keats AS : *Ceiling effect for respiratory depression by nalbuphine. Clin Pharmacol Ther* 27 : 478-485, 1980
- 17) Shin MS, Lee BH, Choi HK, Noh JS, Ahn JY, Sheen SH, et al : *Pain control with continuous infusion of epidural morphine and bupivacaine after lumbar spinal surgery : A prospective study. J Korean Neurosurgery Soc* 29 : 604-608, 2000
- 18) Weis OF, Sriwatanakul K, Alloza JL, Weintraub M, Lasagnal L : *Attitude of patients, house staffs and nurses toward postoperative analgesic care. Anesth Analg* 62 : 70-77, 1983
- 19) Yon SH, Park HC, Kim EY : *The effect of caudal block with morphine sulfate on postoperative pain levels in lumbar spinal surgery. J Korean Neurosurg Soc* 26 : 808-813, 1997